

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/00, 38/04, C07K 1/00, C12N 15/00, 15/64, C12P 21/06		A1	(11) International Publication Number: WO 99/49879
			(43) International Publication Date: 7 October 1999 (07.10.99)
(21) International Application Number: PCT/US99/07189			(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(22) International Filing Date: 31 March 1999 (31.03.99)			
(30) Priority Data: 60/080,083 31 March 1998 (31.03.98) US 09/186,170 4 November 1998 (04.11.98) US			
(71) Applicant: VANDERBILT UNIVERSITY [US/US]; 405 Kirkland Hall, Nashville, TN 37240 (US).			
(72) Inventors: LIN, Yao-Zhong; 576 Harpeth Trace Drive, Nashville, TN 37221 (US). DONAHUE, John, P.; 1601 N. Observatory Drive, Nashville, TN 37215 (US). ROJAS, Mauricio; 3831 Westend Avenue #13, Nashville, TN 37205 (US). TAN, Zhong, Jia; 7420 Stacy Drive, Nashville, TN 37221 (US).			
(74) Agent: LANQUIST, Edward, D., Jr.; Waddey & Patterson, Suite 2020, NationsBank Plaza, 414 Union Street, Nashville, TN 37219 (US).			
(54) Title: SEQUENCE AND METHOD FOR GENETIC ENGINEERING OF PROTEINS WITH CELL MEMBRANE TRANSLOCATING ACTIVITY			
(57) Abstract The invention describes a membrane-translocating peptide sequence (MTS) which facilitates entry of polypeptides and proteins into cells. Also described is an isolated nucleotide sequence encoding the membrane-translocating peptide and a method of using this sequence to genetically engineer proteins with cell membrane permeability. The MTS, and the method of genetically engineering proteins with cell membrane permeability, are useful for polypeptide and protein delivery for human and veterinary applications such as vaccine delivery and cancer therapy.			
BEST AVAILABLE COPY			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SEQUENCE AND METHOD FOR GENETIC ENGINEERING OF
PROTEINS WITH CELL MEMBRANE TRANSLOCATING
ACTIVITY

TECHNICAL FIELD

5 The present invention relates to: (1) DNA sequences encoding membrane-translocating peptide sequences; (2) sequences of membrane-translocating peptides; (3) fusion proteins endowing membrane-translocating potential upon biologically active polypeptides and proteins; and (4) expression vectors for
10 production of said fusion proteins.

 More specifically, the disclosed invention relates to a novel membrane-translocating sequence for directing import of biologically active protein molecules into a cell, and a method of using an expression vector in a host cell to produce a fusion
15 protein comprising a membrane-translocating sequence and a biologically active polypeptide, protein domain, or protein.

BACKGROUND ART

 Signal peptide sequences mediate protein secretion and are composed of a positively charged amino terminus, a central
20 hydrophobic core and a carboxyl-terminal cleavage site recognized by a signal peptidase. These sequences usually comprise 15 to 30 residues. Signal sequences used for targeting proteins to specific locations have been found in both prokaryotic and eukaryotic cells.

 In bacteria, phage fd signal sequences for the major and minor
25 coat proteins direct those proteins to the inner membrane. The -lactamase protein of pBR322 is directed to the periplasmic space by a different signal sequence, while outer membrane proteins such as OmpA are directed to their assigned destination by other signal sequences. Eukaryotic signal sequences directing
30 translocation of the protein into the endoplasmic reticulum include that of human preproinsulin, bovine growth hormone, and the *Drosophila* glue protein. Near the N-terminus of such sequences

are 2-3 polar residues, and within the signal sequence is a hydrophobic core consisting of hydrophobic amino acids. No other conservation of sequence has been observed (Lewin, 1994).

Peptide transport across the cell membrane has been demonstrated, for example, by a peptide representing the third helix of the Antennapedia homeodomain (Derossi *et al.*, 1994). The transport peptide was not used to direct a cargo peptide through the cell membrane, however.

Biological membrane transport has been exploited for protein expression and export from transfected or transformed cells. Secretion of proteins, such as a globin protein, which would normally remain in the cytosol, has been achieved by adding a signal sequence to the N-terminus of the protein (Lewin 1994). Foreign genes have been inserted into recombinant DNA constructs for expression and secretion from bacterial cells, as described for example in U.S. Patent Number 5,156,959, which discloses a method to export gene products into the growth medium of gram negative bacteria. U.S. Patent Number 5,380,653 describes expression vectors and methods for intracellular protein production in *Bacillus* species. U.S. Patent Number 5,712,114 describes a recombinant DNA construct for secretion of expressed proteins, particularly from *Hansenula polymorpha* cells, which utilizes the signal sequence of the human procollagen -1 protein.

Lin *et al.* have described a method of using a naturally-occurring signal peptide sequence to import a cargo peptide into the cell (Lin *et al.*, 1995). One signal sequence that has been successfully used for this cell-permeable peptide import is the 16-residue *h* region of the signal sequence of Kaposi fibroblast growth factor. The cargo peptide transported by this technique has thus far been limited to no more than 25 amino acids, however.

Until now, DNA constructs, including both DNA vaccines

and recombinant viral constructs, have provided the most effective method for furnishing a protein product to the cell for processing and expression of antigenic determinants on the cell surface. The Food and Drug Administration has expressed concern about approval of DNA vaccines, however, citing animal studies in which anti-DNA antibodies have been formed. Recombinant viral vectors have posed a unique set of problems in terms of delivery into cells, efficiency of expression, and potential immune system response to viral proteins. Other methods of DNA transfer into cells, such as transfection and microinjection, are often inefficient and time-consuming.

Genetic disorders resulting from the production of defective protein products have been treated, with limited success, by gene therapy. No other method has shown as much promise for introducing a protein into the interior of a cell. Gene therapy, however, has proven to be more difficult than originally envisioned. Appropriate vectors are difficult to identify, expression is transient, and immune responses to some vectors, particularly viral vectors, may preclude repeated use. Delivery of the isolated protein for import into the affected cells would provide a more efficient and effective solution to the problem.

What is needed is a method for importing entire protein molecules into a cell for studies of intracellular processes in living systems, for drug delivery, for vaccine development, and for disease therapy.

DISCLOSURE OF THE INVENTION

The present invention relates to a novel and non-naturally occurring membrane-translocating sequence (MTS) which has been shown to mediate the transport of a full-length protein into a cell. As used herein, a membrane-translocating sequence is an amino acid sequence capable of mediating the import of a

polypeptide, protein domain, or full-length protein through the cell membrane.

5 The invention further relates to a method of using such a membrane-translocating sequence to genetically engineer proteins with cell membrane permeability. An expression vector is designed so that the DNA sequence encoding the membrane-translocating peptide will be positioned N-terminal or C-terminal to the sequence encoding the target protein, in correct reading frame for expression of both MTS and a biologically-active target protein as a fusion product. Peptides, polypeptides, protein domains, or full-length proteins are expressed as a fusion product with the membrane-translocating sequence.

Expression vectors may be chosen from among those readily available for prokaryotic or eukaryotic expression systems.

15 Genetically engineered proteins prepared by the method of the present invention can be used as protein-based vaccines, particularly where killed or attenuated whole organism vaccines are impractical.

20 Cell-permeable proteins prepared by the method of the present invention can also be used for the treatment of disease, particularly cancer. Cell-permeable proteins can be delivered to the interior of the cell, eliminating the need to transfect or transform the cell with a recombinant vector.

25 Cell permeable polypeptides of the present invention can be used *in vitro* to investigate protein function, or can be used to maintain cells in a desired state.

30 The membrane translocating sequence (MTS) of the present invention can be used to deliver peptides, polypeptides, protein domains, or proteins to the interior of a target cell either *in vitro* or *in vivo*. The MTS can be linked to the target protein through a peptide linkage formed by expression of the fusion protein from a recombinant DNA or RNA molecule, or can be linked to the target

protein by means of a linker covalently linked to the MTS. A covalent linkage can be used to link an MTS of the present invention to a non-protein molecule, such as a polynucleotide, for import into the cell.

5

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the nucleotide and derived amino acid sequences of the membrane-translocating sequence inserted into pGEX-3X to form the expression vectors pGEX-3X-MTS1 (1) and pGEX-3X-MTS2 (2).

10

FIG. 2 illustrates analysis of purified preparations of wild-type glutathione S-transferase (WT-GST), GST-MTS1 fusion protein, and GST-MTS2 fusion protein by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Two micrograms of each protein were separated by 12% SDS-PAGE and stained with Coomassie brilliant blue.

15

FIG. 3a, FIG. 3b, and FIG. 3c. FIG. 3a illustrates indirect immunofluorescence microscopy of NIH 3T3 cells treated with GST-MTS1 fusion protein. FIG. 3b illustrates indirect immunofluorescence microscopy of NIH 3T3 cells treated with GST wild-type protein. FIG. 3c illustrates indirect immunofluorescence microscopy of untreated NIH 3T3 cells.

20

FIG. 4 shows Western blot of cell lysates from NIH 3T3 cells treated with GST-MTS1 or GST-WT. The antibody used for protein detection was anti-glutathione S-transferase.

25

FIG. 5 shows confocal laser scanning microscopy of NIH 3T3 cells treated with 20 M of GST-MTS1 protein for 30 minutes. Protein was detected as yellow fluorescent signals by indirect immunofluorescence assay and analyzed by a six-step Z-position sectional scanning of the cell. Panels 1-7: 1 m cell sections from the bottom to the top of representative GST-MTS1-treated cells. 0: the composite image of all seven sections.

30

FIG. 6a, FIG. 6b, and FIG. 6c. FIG. 6a illustrates concentration-dependence of the cellular import of GST-MTS1 protein. FIG. 6b illustrates temperature-dependence of the cellular import of GST-MTS1 protein. FIG. 6c illustrates time-dependence of the cellular import of GST-MTS1 protein. As shown in FIG. 6a, cells were treated with 0-20 mM concentrations of GST-MTS1 protein, as indicated. Total cell lysates were then analyzed by Western blot analysis using anti-GST antibody as probe. As shown in FIG. 6b, cells were treated with equal concentrations of GST-MTS1 protein at 4°C, 22°C, and 37°C, as indicated. Total cell lysates were then analyzed by Western blot with anti-GST antibody as probe. As shown in FIG. 6c, GST-MTS1 continued to accumulate intracellularly up to 18 hours of incubation, with only a small percentage of imported protein appearing degraded. Cells treated with 20 M protein at 37°C for the times indicated continued to import the protein.

FIG. 7a and FIG. 7b. FIG. 7a shows indirect immunofluorescence microscopy of serum-starved SAA cells treated with 2.5 mM GST-Grb2SH2-MTS. FIG. 7b shows indirect immunofluorescence microscopy of serum-starved SAA cells treated with 2.5 mM GST-Grb2SH2 protein for one hour, followed by treatment with epidermal growth factor (EGF) (50 ng/ml) for 10 minutes.

FIG. 8a and FIG. 8b. FIG. 8a illustrates Western blot analysis of cell lysates from SAA cells treated with the indicated proteins and EGF. Probes were specific for phosphorylated EGF receptor (top panel) and Grb2 protein (bottom panel). FIG. 8b illustrates Western blot analysis of anti-Grb2 immunoprecipitates of cell lysates from SAA cells treated with GST-Grb2SH2-MTS at the indicated concentrations followed by EGF treatment (50 ng/ml) for 10 minutes.

FIG. 9 illustrates Western blot analysis, using anti-GST

antibody, of the intracellular level of GST-Stat1SH2 imported into the cell as a GST-Stat1SH2-MTS fusion product. Lysates of untreated cells were run in the lane marked "-", and lysates of cells treated with the GST-Stat1SH2-MTS fusion product were run in the lane marked "GST-Stat1SH2-MTS." The GST-Stat1SH2-MTS product migrated to the predicted 43kDa location, demonstrating that a protein of this size could be imported into the cell when fused to the MTS.

10 BEST MODE FOR CARRYING OUT THE INVENTION

Previous efforts to import small peptides into a cell have been successful, but efforts to import larger polypeptides or whole proteins have not. The present invention seeks to overcome the shortcomings of the prior art by providing a peptide capable of directing the import of polypeptides and proteins into a cell. It is to be understood that, as used herein, polypeptide is intended to encompass any amino acid sequence comprising more than three amino acids, and includes particularly protein domains and proteins.

20 The present invention relates to a membrane-translocating peptide and its use in mediating membrane-translocation and import of a polypeptide, protein domain, or full-length protein into a cell. The inventors have synthesized an artificial membrane translocation sequence (MTS) of 12 amino acids for protein import, and have used a DNA sequence encoding this 12-amino acid peptide to construct a plasmid expression vector for genetically engineering proteins with cell membrane permeability by expressing the 12 amino acid MTS as a fusion with a target protein for import into the cell. In the preferred embodiment of the present invention, the amino acid sequence of the 12-residue membrane-translocating peptide is Ala-Ala-Val-Leu-Leu-Pro-Val-Leu-Leu-Ala-Ala-Pro (SEQ. ID NO. 1). As used herein, however,

the term "peptide" is intended to include mimetics and the term "amino acid" is intended to include D-form amino acids and modified amino acids. These substitutions may be made by someone of skill in the art, using the known structural similarities between the molecules. The membrane translocating sequence may be located immediately adjacent to, or some distance from, the cargo protein as produced by the recombinant nucleotide vector of the present invention. Therefore, the amino acid sequence is also intended to include any peptide or protein sequence that may include additional amino acids either N-terminal or C-terminal to the listed sequence, or both.

The amino acid sequence is also intended to include an MTS comprising fewer than twelve residues, as signal peptide sequences of as few as eight amino acids provide membrane translocation of peptides across membranes within the cell. In the present invention, an alternative MTS is comprised of an amino acid sequence of eight (8) to twelve (12) consecutive amino acids chosen from SEQ. ID NO. 1. Exemplary of such alternative MTS sequences are Ala-Ala-Val-Leu-Leu-Pro-Val-Leu (SEQ. ID NO. 2), Ala-Ala-Val-Leu-Leu-Pro-Val-Leu-Leu (SEQ. ID NO. 3), Ala-Ala-Val-Leu-Leu-Pro-Val-Leu-Leu-Ala (SEQ. ID NO. 4), Ala-Ala-Val-Leu-Leu-Pro-Val-Leu-Leu-Ala-Ala (SEQ. ID NO. 5), Leu-Pro-Val-Leu-Leu-Ala-Ala-Pro (SEQ. ID NO. 6), Leu-Leu-Pro-Val-Leu-Leu-Ala-Ala-Pro (SEQ. ID NO. 7), Val-Leu-Leu-Pro-Val-Leu-Leu-Ala-Ala-Pro (SEQ. ID NO. 8), and Ala-Val-Leu-Leu-Pro-Val-Leu-Leu-Ala-Ala-Pro (SEQ. ID NO. 9). Alternative MTS sequences are intended to include alternative amino acids, as well as additional C-terminal or N-terminal amino acids as described for SEQ. ID. NO. 1.

In a second preferred embodiment of the invention, the DNA coding sequence for the membrane-translocating peptide is 5'-GCAGCCGTT CTTCTCCCTGTTCTTCTTGCCGCACCC-3'

(SEQ. ID NO. 10). Alternate embodiments include, but are not limited to: 5'-GCAGCCGTT CTTCTCCCTGTTCTT-3' (SEQ. ID NO. 11); 5'-GCAGCCGTT CTTCTCCCTGTTCTTCTT-3' (SEQ. ID NO. 12); 5'-GCAGCCGTT CTTCTCCCTGTTCTTCTTGCC-3' (SEQ. ID NO. 13); 5'-GCAGCCGTT CTTCTCCCTGTTCTTCTTGCCGCA-3' (SEQ. ID NO. 14); 5'-CTCCCTGTTCTTCTTGCCGCACCC-3' (SEQ. ID NO. 15); 5'-CTTCTCCCTGTTCTTCTTGCCGCACCC-3' (SEQ. ID NO. 16); 5'-GTTCTTCTCCCTGTTCTTCTTGCCGCACCC-3' (SEQ. ID NO. 17); and 5'-GCCGTT CTTCTCCCTGTTCTTCTTGCCGCACCC-3' (SEQ. ID NO. 18). It is well known in the art, however, that a single amino acid may be encoded by more than one nucleotide codon—and that the nucleotide sequence may be easily modified to produce an alternate nucleotide sequence that encodes the same peptide. Therefore, alternate embodiments of the present invention include alternate DNA sequences encoding peptides containing the amino acid sequences as previously described. DNA sequences encoding peptides containing the claimed amino acid sequence include DNA sequences which encode any combination of the claimed sequence and any other amino acids located N-terminal or C-terminal to the claimed amino acid sequence.

It is to be understood that amino acid and nucleic acid sequences may include additional residues, particularly N- or C-terminal amino acids or 5' or 3' nucleotide sequences, and still be essentially as set forth in the sequences disclosed herein, as long as the sequence confers membrane permeability upon the polypeptide or protein moiety of the fusion protein.

A nucleic acid fragment of almost any length may be employed, and may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like. Therefore, overall length may vary considerably.

In the method of the present invention, the nucleotide sequence described above is inserted into a protein expression vector to produce a protein which can be imported from the exterior to the interior of a cell by the action of the membrane translocating sequence described herein.

Genetically Engineering a Protein with Cell Membrane Permeability

In a preferred embodiment of the present invention, a protein expression vector is genetically engineered to incorporate a DNA sequence encoding a membrane translocating sequence in an orientation either N-terminal or C-terminal to the DNA sequence encoding the peptide, polypeptide, protein domain, or full-length protein of interest, and in correct reading frame so that a fusion protein consisting of the membrane translocating peptide and the target peptide, polypeptide, protein domain, or full-length protein may be expressed. It is understood by those of skill in the art that all protein domains and full-length proteins are polypeptides, being formed of a series of peptide linkages. Therefore, as used herein, the term "polypeptide" will be used to denote an amino acid sequence of more than 25 amino acids, and the term "peptide" will be used to denote an amino acid sequence of 25 amino acids or less. In the preferred embodiment of the method of genetically-engineering proteins with cell membrane permeability as described in the present invention, the membrane-translocating peptide is an MTS of the present invention. In alternate embodiments of the present method of genetically engineering proteins with membrane translocating activity, the MTS may comprise an alternate sequence which mediates the import of a peptide or polypeptide through the cell membrane to the interior of a cell.

In a further embodiment of the invention, a cleavage site is

located between the MTS and the target polypeptide, protein domain, or full-length protein. This site may alternatively be a factor X site, or other site that is known to those of skill in the art to effect the cleavage of the fusion protein to physically remove the
5 MTS from the subject peptide or polypeptide. As used herein, an MTS is a membrane translocating sequence of the present invention, which directs cellular transport of a target protein from the exterior to the interior of a cell. A target protein is a protein which normally evidences less than optimal permeability through
10 the cell membrane, but which, when linked either N-terminal or C-terminal to an MTS of the present invention, is transported from the exterior to the interior of the cell.

The MTS of the present invention, and the method of genetically engineering proteins for cell membrane permeability,
15 can be used in a variety of applications, including, but not limited to, studies of intracellular protein function, vaccine delivery, and delivery of peptides, nucleic acids, and other organic compounds for therapeutic use. A specific example of a polypeptide involved in intracellular signaling is the SH2 domain of the Grb2 protein that
20 becomes bound to tyrosine-phosphorylated epidermal growth factor receptor (EGFR). A specific example of a viral protein is the Hepatitis B surface antigen, or the human immunodeficiency virus type 1 HXB-2 envelope glycoprotein. The MTS of the present invention has been shown thus far to mediate the cellular import
25 of an entire 120-kDa protein fusion product. Other proteins that can be delivered to the interior of the cell using the method of the present invention include, but are not limited to, MAP kinase, RAS, caspases, protein members of the Bcl-2 family, Bax, NF B, green fluorescent protein (GFP) and STAT.

30 The method of the present invention provides a means of producing proteins with cell permeability for introduction into the interior of the cell, where their actions help to further elucidate

cellular control and biosynthesis mechanisms. This method also provides a means to introduce intracellular proteins into cells to produce targeted cellular changes, such as inhibition of apoptosis by the introduction of Bcl-2. Cell cycle control, for example, can be
5 altered by the introduction of a functional p53 protein product into those cells that have become tumorigenic due to an abnormal p53 protein.

Expression system vectors, which incorporate the necessary regulatory elements for protein expression, as well as restriction
10 endonuclease sites that facilitate cloning of the desired sequences into the vector, are known to those of skill in the art. A number of these expression vectors are commercially available. In a preferred embodiment of the present invention, the expression vector is pGEX-3X (Amersham Pharmacia, Piscataway NJ), U.S.
15 Patent Number 5,654,176 (Smith, *et al.*, incorporated herein by reference), which comprises a nucleotide sequence encoding a fusion protein including glutathione-S-transferase. Insertion of a nucleotide sequence encoding an MTS as described by the present invention into vector pGEX-3x, either 5' or 3' to the glutathione-S-
20 transferase (GST) gene of the vector enables expression of a fusion protein incorporating both the MTS and the glutathione-S-transferase protein. The MTS, connected either N-terminal or C-terminal to the glutathione-S-transferase protein, carries the GST protein through the cell membrane to the interior of the cell.

25 In another preferred embodiment of the present invention, an alternate recombinant DNA expression vector containing the elements previously described is introduced into an appropriate host cell where cellular mechanisms of the host cell direct the expression of the fusion protein encoded by the recombinant DNA
30 expression vector. Alternately, cell-free systems known to those of skill in the art can be chosen for expression of the fusion protein.

The purified fusion protein produced by the expression

vector host cell system can then be administered to the target cell, where the membrane-translocating sequence mediates the import of the fusion protein through the cell membrane of the target cell into the interior of the cell.

5 An expression vector host cell system can be chosen from among a number of such systems that are known to those of skill in the art. In one embodiment of the invention, the fusion protein can be expressed in *Escherichia coli*. In alternate embodiments of the present invention, fusion proteins may be expressed in other
10 bacterial expression systems, viral expression systems, eukaryotic expression systems, or cell-free expression systems. Cellular hosts used by those of skill in the art include, but are not limited to, *Bacillus subtilis*, yeast such as *Saccharomyces cerevisiae*, *Saccharomyces carlsbergensis*, *Saccharomyces pombe*, and *Pichia*
15

pastoris, as well as mammalian cells such as 3T3, HeLa, and Vero. The expression vector chosen by one of skill in the art will include promoter elements and other regulatory elements appropriate for the host cell or cell-free system in which the fusion protein will be expressed. In mammalian expression systems, for example,
20 suitable expression vectors can include DNA plasmids, DNA viruses, and RNA viruses. In bacterial expression systems, suitable vectors can include plasmid DNA and bacteriophage vectors.

 Examples of specific expression vector systems include the
25 pBAD/gIII vector (Invitrogen, Carlsbad, CA) system for protein expression in *E. coli*, which is regulated by the transcriptional regulator AraC. Dose-dependent induction enables identification of optimal expression conditions for the specific target protein to be expressed. By inserting the polynucleotide sequence of the
30 membrane translocating sequence of the present invention either 5' or 3' to the polynucleotide sequence of a target protein, this vector can be used to express a number of fusion proteins for which

optimal expression conditions may vary. Furthermore, the vector encodes the polyhistidine (6xHis) sequence and an epitope tag to allow rapid purification of the fusion protein with a nickel-chelating resin, along with protein detection with specific antibodies to detect the presence of the secreted protein.

An example of a vector for mammalian expression is the pcDNA3.1/V5-His-TOPO eukaryotic expression vector (Invitrogen). In this vector, the fusion protein can be expressed at high levels under the control of a strong cytomegalovirus (CMV) promoter. A C-terminal polyhistidine (6xHis) tag enables fusion protein purification using nickel-chelating resin. Secreted protein produced by this vector can be detected using an anti-His (C-term) antibody.

Another example of a protein expression vector for a mammalian expression system is the pEBVHis (Invitrogen) vector. There are three different versions of this vector (pEBVHis A, pEBVHis B, and pEBVHis C), that differ in spacing between the sequences that code for the N-terminal peptide and the multiple cloning site. The vector can therefore be chosen to facilitate cloning the target polypeptide nucleotide sequence into the vector in correct reading frame to produce a biologically functional polypeptide. The multiple cloning has nine unique restriction sites, including BamHI, XhoI, Bgl II, Pvu II, Kpn I, Hind III, Not I, Sfi I, and Cla I to facilitate insertion of the nucleotide sequences for production of the MTS/target polypeptide fusion. The vector includes an Epstein-Barr virus origin of replication, and the Epstein-Barr virus encoded nuclear antigen (EBNA-1) which transactivates the origin of replication to allow the vector DNA to replicate episomally when transfected into an appropriate mammalian cell line. Appropriate cell lines for this vector include 293 cells, 293-EBNA, COS, or CV-1. The Rous Sarcoma virus long terminal repeat directs transcription of the fusion protein in this

vector, and selection in mammalian cells is facilitated by the Hygromycin B drug resistance marker under the control of the thymidine kinase promoter in the vector. Purification of the fusion protein can be accomplished using metal affinity chromatography to bind the polyhistidine tag, and the tag can be subsequently
5 cleaved from the fusion protein using an enterokinase cleavage recognition sequence.

A baculovirus expression system can also be used for production of a fusion protein comprising the MTS and a target
10 protein. A commonly used baculovirus is AcMNPV. Cloning of the MTS/target protein DNA can be accomplished by using homologous recombination. The MTS/target protein DNA sequence is cloned into a transfer vector containing a baculovirus promoter flanked by baculovirus DNA, particularly DNA from the polyhedrin gene. This DNA is transfected into insect cells, where
15 homologous recombination occurs to insert the MTS/target protein into the genome of the parent virus. Recombinants are identified by altered plaque morphology.

Many fusion proteins containing target proteins that may
20 not be appropriately post-translationally modified in bacterial expression systems can be expressed with baculovirus vectors. In a method for generating recombinant baculovirus, the MTS/target protein DNA is cloned into donor plasmid, such as the pFastBac donor plasmid of the Bac-to-Bac™ Baculovirus Expression System
25 (GibcoBRL). The recombinant plasmid is then transformed into *E. coli* host cells that contain a bacmid with a mini-*att*Tn7 target site and helper plasmid. The mini-Tn7 element on the donor plasmid can transpose to the mini-*att*Tn7 target site on the bacmid in the presence of helper plasmid transposition proteins. Transposition
30 results in disruption of the *lacZ* gene, allowing identification of colonies containing recombinant bacmids. Recombinant bacmid DNA is then used to transfect insect cells, such as *Spodoptera*

frugiperda cell Sf9. By using a histidine tag expression vector, such as the pFastBacHT expression vector (Gibco BRL), the expressed fusion protein can be purified using the 6xHis tag.

5 Enzymes, signaling molecules, mediators of cell cycle control, transcription factors, antigenic peptides, full-length protein products of viral, bacterial, or other origin for use in vaccine therapy, protein products of human cells for use in cancer vaccine therapy, toxins, and proteins involved in intracellular signaling systems which may not be appropriately post-
10 translationally modified in bacterial expression systems can be expressed with baculovirus vectors.

Proteins as described above can also be produced in the method of the present invention by mammalian viral expression systems. The Sindbis viral expression system, for example, can be
15 used to express the fusion protein at high levels. Sindbis vectors have been described, for example, in U.S. Patent Number 5,091,309 (Schlesinger *et al.*), incorporated herein by reference. Sindbis expression vectors, such as pSinHis (Invitrogen, Carlsbad, CA) can be used to express the fusion protein under the direction
20 of the subgenomic promoter PSG. *In vitro* transcribed RNA molecules encoding the fusion protein and the Sindbis proteins required for *in vivo* RNA amplification can be electroporated into baby hamster kidney (BHK) cells using methods known to those of skill in the art. Alternatively, the RNA encoding the fusion
25 protein and Sindbis proteins required for *in vivo* RNA amplification can be cotransfected with helper RNA that permits the production of recombinant viral particles. Viral particles containing genetic material encoding the fusion protein can then be used to infect cells of a wide variety of cell types, including
30 mammalian, avian, reptilian, and *Drosophila*. Fusion protein expressed from the pSinHis (Invitrogen) vector can be detected with antibody to an Anti-Xpress™ epitope encoded by the vector

sequence. The pSinHis vector also includes a polyhistidine tag which provides a binding site for metal-chelating resins to facilitate purification of the expressed fusion protein. Furthermore, an enterokinase cleavage site located between the histidine tag and the fusion protein allows the histidine tag to be enzymatically removed following purification.

An ecdysone-inducible mammalian expression system (Invitrogen, Carlsbad, CA), described by No, *et al.* (1996) can also be used to express the MTS/target fusion protein. Vectors used in the ecdysone-inducible mammalian expression system can be organized to produce cell-permeable target proteins by expressing the MTS/target fusion protein from the expression cassette. With the ecdysone-inducible system, higher levels of protein production can be achieved by use of the insect hormone 20-OH ecdysone to activate gene expression via the ecdysone receptor. An inducible expression plasmid provides a multiple cloning site, into which the nucleotide sequence of the MTS and a target protein can be ligated, oriented so that the MTS is translated either N-terminal or C-terminal to the target protein in the expressed fusion protein. The expression vector contains ecdysone response elements upstream of the promoter (a minimal heat shock promoter) and the multiple cloning site. Cotransfection of a second plasmid, pVgRXR (Invitrogen), provides the receptor subunits to make the cell responsive to the steroid hormone ecdysone analog, ponasterone A. A control expression plasmid containing the *lacZ* gene can be cotransfected with pVgRXR to provide a marker for transfected cells. Upon induction with ponasterone A, the control plasmid expresses -galactosidase. Cotransfection of the inducible expression construct and pVgRXR into the mammalian cell of choice can be accomplished by any of the standard means known to those of skill in the art. These include, for example, calcium phosphate transfection, lipid-mediated transfection, and

electroporation. Levels of expression of the fusion protein in this system can be varied according to the concentration and length of exposure to ponasterone. Stable cell lines that constitutively express the MTS/target fusion protein can be established using
5 Zeocin™ (Invitrogen), a bleomycin/phleomycin-type antibiotic isolated from *Streptomyces*, and neomycin or hygromycin.

Yeast host cells, such as *Pichia pastoris*, can also be used for the production of a genetically engineered cell permeable protein by the method of the present invention. Expression of
10 heterologous proteins from plasmids transformed into *Pichia* has previously been described by Sreekrishna, *et al.* (U.S. Patent Number 5,002,876, incorporated herein by reference). Vectors for expression in *Pichia* of a fusion protein comprising an MTS of the present invention and a target protein are commercially available
15 as part of a *Pichia* Expression Kit (Invitrogen, Carlsbad, CA). *Pichia pastoris* is a methylotrophic yeast, which produces large amounts of alcohol oxidase to avoid the toxicity of hydrogen peroxide produced as a result of methanol metabolism. Alcohol oxidase gene expression is tightly regulated by the AOX1 and
20 AOX2 promoters. In *Pichia* expression vectors, high levels of expression are produced under the control of these promoters. Ohi, *et al.* (U.S. Patent Number 5,683,893, incorporated herein by reference) have previously described a mutant AOX2 promoter capable of producing enhanced expression levels. Using previously
25 described and commercially available *Pichia* expression vectors, a target protein can be genetically engineered for cell permeability by incorporating into the expression vector both the nucleotide sequence of the target protein and a nucleotide sequence encoding an MTS of the present invention. The nucleotide sequence
30 encoding an MTS can be incorporated into the vector either 5' or 3' to the nucleotide sequence encoding the target protein. Under the control of the AOX1 or AOX2 promoter, high levels of protein can

be expressed.

Purification of heterologous protein produced in *Pichia* has been described by Craig, *et al.* (U.S. Patent Number 5,004,688, incorporated herein by reference), and techniques for protein
5 purification from yeast expression systems are well known to those of skill in the art. In the *Pichia* system, commercially available vectors can be chosen from among those that are more suited for the production of cytosolic, non-glycosylated proteins and those that are more suited for the production of secreted, glycosylated
10 proteins, or those directed to an intracellular organelle, so that appropriate protein expression can be optimized for the target protein of choice.

Peptide Attachment Using Covalent Linkage

15 The MTS of the present can also be used to increase cell membrane permeability of a polypeptide, oligonucleotide, or other organic molecule by attaching the MTS to the target molecule by means of a covalent attachment. Orthogonal coupling methods for peptides and polypeptides involving a thioester intermediate have
20 been described by Tam, *et al.* (1995).

Therapeutic use of oligonucleotides is often hindered by their low cellular permeability. Although oligonucleotides have been shown to be taken up by cells via an endocytic process, oligonucleotides that enter the cell in this manner are usually
25 trapped in endocytic vesicles and degraded in lysosomes. Dokka, *et al.* (1997) have demonstrated non-endocytic uptake of oligonucleotides using a signal import peptide consisting of the hydrophobic sequence of Kaposi fibroblast growth factor signal peptide (Ala-Ala-Val-Ala-Leu-Leu-Pro-Ala-Val-Leu-Leu-Ala-Leu-
30 Leu-Ala-Pro) covalently conjugated to a polycationic linker, poly-L-lysine (PL). The poly-L-lysine linker was further complexed electrostatically to the polyanionic backbone of the oligonucleotide.

The signal peptide/poly-L-lysine linker was synthesized using the standard Fmoc procedure.

5 The MTS of the present invention provides efficient transport of large protein molecules across the cell membrane, whereas other membrane transport peptides previously tested have not been demonstrated to transport molecules of size greater than approximately 25 amino acids. An MTS as described herein can be attached to a peptide or polypeptide using methods, such as those described by Tam *et al.*, to enhance peptide or polypeptide membrane permeability. The MTS can be provided in the form of a kit, including the necessary components known to those of skill in the art to facilitate linkage of a peptide to a target polypeptide. A target protein linked to the MTS in this manner can then be delivered to the cell either *in vitro* or *in vivo* for intracellular import.

15 An MTS of the present invention can also be provided as a fusion between the MTS and a poly-L-lysine linker. When this fusion product is mixed with oligonucleotide, as described by Dokka, et al., the poly-L-lysine linker can complex electrostatically with the polyanionic backbone of the oligonucleotide. The MTS-PL linker-oligonucleotide complex can then be administered *in vitro* or *in vivo* to deliver the oligonucleotide to the interior of the cells.

Conditions for Protein Import Into a Target Cell

25 Suitable conditions for protein import into the cell mediated by the membrane-translocating peptide of the present invention include incubating the cells in an extracellular concentration of fusion protein in the 20 M range at 37°C for 30 minutes, to accomplish the import of approximately $0.5-1 \times 10^6$ molecules of transported protein per cell. Effective concentrations, however, may vary with differing proteins and cell types, and may be considered as amounts sufficient to result in import of fusion

30

proteins into the cell, with protein import exhibiting dose-dependence. Methods for providing sufficient concentration to achieve protein import are known to those of skill in the art. Suitable import temperatures include temperatures in a preferred
5 range between 22°C and 37°C.

The fusion protein produced by the method of the present invention may be administered *in vitro* by any of the standard methods known to those of skill in the art, such as addition of fusion protein to culture medium, or other methods as described by
10 Lin *et al.*, U.S. Patent No. 5,807,746, incorporated herein by reference. Furthermore, it will be appreciated by those of skill in the art that fusion proteins produced by this method may be delivered *in vivo* by standard methods utilized for protein/drug delivery, including parenteral administration, intravenous
15 administration, topical administration, aerosol administration or inhalation, oral administration (particularly when provided in encapsulated form), or by rectal or vaginal administration (particularly when provided in suppository form).

Administration of fusion protein produced by the method of the present invention may be performed for a time length of from
20 30 minutes to 18 hours, particularly when administration is accomplished by addition of fusion protein to culture media for *in vitro* use. For *in vivo* or *in vitro* use, effective administration time for a fusion protein produced by the method of the present invention may be readily determined by one of skill in the relevant
25 art.

Uptake of the fusion protein is dependent upon the external concentration of the fusion protein and the period of application, therefore the internal concentration of protein can be controlled by
30 controlling administration to the extracellular environment. At an extracellular concentration of 20 M, NIH3T3 cells, for example, receive up to 10^6 molecules of protein per cell over a 60-minute

period, resulting in an internal concentration of approximately 2 M.

Studying Intracellular Proteins Using MTS and Molecular Labeling

5 Molecular labeling techniques have previously been described for studying intracellular protein function. Jones, *et al.* (1998) describe a method of molecular labeling using intracellular expression of a fusion protein comprising green fluorescent protein (GFP) and dynamin. Kohler, *et al.* (1997) describe a method of
10 studying exchange of protein molecules through the connections between plastids of higher plants. The molecular exchange was visualized using green fluorescent protein (GFP) to label the plastid stroma. In fact, GFP from *Aequorea victoria* has been
15 genetically fused with many host proteins to produce fluorescent chimeras (reviewed by Tsien, 1998). Griffin, *et al.* (1998) describe a method of incorporating a small receptor domain, utilizing four cysteines at the i, i+1, i+4, and i+5 positions of an α -helix, to create a ligand for 4',5'-bis(1,3,2-dithioarsolan-2-yl)fluorescein,
20 which is membrane-permeant and non-fluorescent until it binds with high affinity and specificity to the tetracysteine domain. This *in situ* labeling technique provides greater versatility in attachment sites and eliminates the potentially disruptive effects of GFP (a 238 amino acid protein) in the cell.

25 Olson, *et al.* (1995) describe a method of analyzing MAP4 function in living cells using a GFP-MAP4 chimera. Expression of the GFP-MAP4 chimera in dividing cells enabled visualization of MAP4 in microtubule organization. Rizzuto, *et al.* (1995) describe a modified GFP cDNA that includes a mitochondrial targeting
30 sequence. Expression of the cDNA construct in the cells allows visualization of mitochondrial movement within living cells.

The techniques previously described, however, rely upon

protein expression within the target cell using polynucleotide vectors. The MTS of the present invention, and the method of genetically engineering proteins for membrane permeability, provides a more efficient method for studying intracellular protein function by producing a fusion protein comprising the MTS, a protein label (such as GFP of the tetracysteine domain described by Griffin), and the protein to be studied. The fusion protein can be produced in a system such as those described previously, and the purified fusion protein can be administered to the cells. Once administered to the extracellular environment, the MTS directs import of the chimeric protein into the interior of the cell and the molecular marker enables visualization of target protein localization.

Other markers that may be used in the method of the present invention include, but are not limited to, rhodamine, biotinylated markers, and blue fluorescent protein. Vector systems providing the polynucleotide coding sequence for green or blue fluorescent proteins are available from Aurora Biosciences (San Diego, CA), Clontech (Palo Alto, CA), and Quantum Biotechnologies (Montreal, Canada). Multiple cloning sites within these vectors enable insertion of the MTS of the present invention either N-terminal or C-terminal to the label/target protein chimera.

The method previously described can also be used to label cells to facilitate observation of, for example, cellular migration through tissue and tumor metastasis. An MTS/GFP/target protein chimera can be administered to cells prior to injection *in situ*, or can be administered locally *in situ*, in the case of tumor cells, to study metastasis. Methods for studying cellular migration are known to those of skill in the art. These methods are facilitated by the method of the present invention, producing a chimeric protein which labels the cell with a protein that is easily detectable, cell-

permeable, and therefore located in the interior of the cell due to the presence of the MTS either C-terminal or N-terminal to the GFP or target protein sequence.

5 ***Vaccine Administration Using Membrane Translocating Sequences and Genetically Engineered Proteins***

Vaccines provide the most effective means of control of infectious disease. Deaths from infectious diseases rose 22 percent between 1980 and 1992 (not including deaths attributable to HIV; the virus which causes acquired immunodeficiency syndrome). In an effort to reduce the prevalence of infectious disease and decrease the risks associated with vaccination with killed or attenuated live organisms, scientists have focused on the development of peptide vaccines and DNA vaccines. Organisms for which antigenic proteins have been identified include *Haemophilus influenzae* B, *Clostridium difficile*, *Helicobacter pylori*, meningococcus, and *Borrelia burgdorferi*, to name only a few. For example, the 31 kD antigen OspA and 34 kD antigen OspB of *Borrelia burgdorferi* have been demonstrated to provide a sufficient protective response to provide the incentive to begin the required FDA vaccine trials for a lyme disease vaccine using these protein antigens. Viruses for which vaccines are currently being developed include human immunodeficiency virus (HIV), Ebola, influenza, cytomegalovirus, Epstein-Barr virus, herpes simplex, human papillomavirus, parainfluenza type 3, and B19 parvovirus. Recombinant vaccines, often utilizing viral vectors, have provided the most promising vector for antigen delivery. However, administration requires administration of the vector rather than just the protein product. The viral vectors have proven to present safety issues which often limit their use.

Viral vaccines, although effective, often pose serious problems in terms of delivery and in terms of control of transmission. Foot-and-mouth disease virus (FMDV), for example, is a highly contagious viral disease of pigs and cattle. Inactivated virus vaccines are effective, but outbreaks of the disease have been directly associated with incomplete inactivation of virus or the escape of virus from vaccine manufacturing facilities (King *et al.*, 1981). Efforts have therefore been directed toward the development of DNA vaccines, which produce isolated proteins of the infectious agent and eliminate the possibility of reversion to virulence. Chinsamgaram, *et al.* (1998) developed a DNA inoculation-based strategy to utilize plasmid DNA to produce non-infectious viral capsids in inoculated animals. Inoculated animals subsequently developed antibodies to the capsid proteins, providing a protective response. Chen, *et al.* (1998) demonstrated that PLG-encapsulated rotavirus VP6 DNA produced a protective response after oral administration in BALB/c mice.

U.S. Patent No. 5,703,057, issued to Johnston *et al.* (incorporated herein by reference), describes the use of vaccines based upon expression libraries constructed from fragmented genomic DNA of pathogens. Once transfected into the host cell, the proteins derived from the DNA sequences are produced. Advantages of this technique include: (1) production of vaccines without having to predetermine which specific proteins are responsible for eliciting protection; (2) presentation of peptides which might normally be hidden by immune-avoidance mechanisms in the killed or attenuated host organism; (3) presentation of whole protein antigens in a manner similar to that achieved by live/attenuated vaccines; (4) modification of the vaccine composition to utilize only those antigenic proteins found to be most effective; and (5) introduction of antigens into cells which might normally not be affected by live/attenuated

organisms. These same advantages are inherent in the present invention. Peptide vaccines of the present invention, however, provide the further advantage of providing a polypeptide or protein to the cell without requiring administration of a recombinant vector, which may or may not provide efficient protein production.

Peptide delivery has been accomplished by a variety of means, including encapsulation, and has been demonstrated to produce a sufficient antigenic response to provide protection. Peptide vaccines, however, require identification of specific antigenic epitopes in order to synthesize the appropriate peptide for vaccine use. Immunization with a complete protein or protein domain provides a method for introducing epitopes into the cell without having to first isolate the peptides containing them. The antigen-processing machinery of the cell then provides the antigen processing necessary to invoke a protective immune response.

Synthetic peptides from the envelope glycoprotein sequence of Murray Valley encephalitis (MVE) virus have been shown by Mathews, *et al.* (1992) to induce antibody and *in vitro* proliferation of peptide-primed helper T (Th) cells. In dogs, a synthetic peptide vaccine has been demonstrated to protect dogs against challenge with virulent canine parvovirus (Langeveld, *et al.* 1994).

Methods for preparation of vaccines containing peptide sequences as active ingredients are well known in the art. Such methods are exemplified in U.S. Pat. Nos. 4,578,770; 4,596,792; 4,599,230; 4,599,231; 4,608,251; and 4,601,903, all incorporated herein by reference. These methods can also be used to prepare vaccines using cell-permeable polypeptides and proteins.

Vaccines using cell-permeable polypeptides provide an advantage over peptide vaccines. Immune system recognition of antigen depends upon appropriate antigen processing. Previously, entire proteins or protein domains could not be delivered to the

interior of the cell for processing to occur. As a consequence, peptides representing antigenic epitopes had to be identified prior to delivery to the cell of small peptides representing those epitopes. The method of the present invention allows whole
5 proteins or protein domains to be imported into the cell, where antigenic processing can occur. This provides multiple antigenic epitopes in one administration, and eliminates the need for experimental identification of specific epitopes for vaccine development.

10 Typically, such vaccines are prepared for injection into a human or mammalian subject. Injectable vaccines can be prepared as liquid solutions or suspensions. Solid forms can be prepared which are suitable for solution in, or suspension in, liquid prior to injection. The preparation may also be emulsified.
15 The active immunogenic ingredient is often mixed with a pharmaceutically acceptable carrier which is compatible with the active ingredient. Suitable carriers include, but are not limited to, water, dextrose, glycerol, saline, ethanol, and combinations thereof. The vaccine may contain additional agents such as
20 wetting or emulsifying agents, pH buffering agents, or adjuvants which enhance the effectiveness of the vaccine.

The vaccine may be conventionally administered parenterally. Either subcutaneous or intramuscular injection is appropriate. Other modes of administration may include oral
25 administration, nasal administration, rectal administration, and vaginal administration, which may involve combining the peptide immunogen with pharmaceutically acceptable carriers such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, or other carrier. Compositions for
30 oral administration may form solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders. A protein-based vaccine of the present invention can be administered by

enteric-coated capsule for release of the polypeptide into the lumen of the intestine.

The peptides of the present invention may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, mandelic, oxalic, and tartaric. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, and histidine.

The vaccine is administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, taking into account, for example, the capacity of the individual's immune system to synthesize antibodies, and the degree of protection desired. Precise amounts of active ingredient (peptide immunogen) to be administered depend on the judgment of the practitioner. Suitable dosage ranges generally require several hundred micrograms of active ingredient per vaccination. Also variable are regimes for initial administration and booster vaccinations, which should be determined by the judgment of the practitioner. Dosage of vaccine will depend on the route of administration and will vary according to the size of the host.

Adjuvants for use in combination with the polypeptide immunogen of the present invention for vaccination include, but are not limited to, aluminum hydroxide or phosphate, also known as alum, commonly used as 0.05 to 0.1 percent solution; aggregation of the protein in the vaccine by heat treatment with

temperatures ranging between 70° for 30 seconds to 101° C for 2 minutes.

Methods for producing the hepatitis B surface antigen (HBsAg) in yeast have been described previously in U. S. Patent
5 No. 4,769,238 issued to Rutter, *et al.*, U.S. Patent No. 4,895,800 issued to Tschopp, *et al.*, and U.S. Patent No. 5,098,704 issued to Valenzuela (all incorporated herein by reference). Using the method of the present invention, HBsAg can be produced in yeast, such as *Pichia pastoris*, as a fusion protein containing the
10 membrane translocating sequence (MTS) either N-terminal or C-terminal to the HbsAg protein. The purified fusion product can then be administered as a protein vaccine, capable of entering the cell for antigen-processing by means of the membrane translocating sequence.

15 Xu, *et al.* have described a method of immunizing guinea pigs with plasmids encoding viral glycoproteins from the Ebola virus. A vaccine delivered by their method has been demonstrated to provide protective immunity, and protection correlated with antibody titer and antigen-specific T-cell responses to the viral
20 glycoproteins. In the method of the present invention, the virion glycoproteins, both secreted form (sGP) and transmembrane form (GP), of Ebola can be produced in a suitable vector, such as *Pichia pastoris*, as a fusion protein with the MTS located either N-terminal or C-terminal to the virion glycoprotein. The purified
25 fusion protein can be delivered by standard means of vaccination for peptide or subunit-based vaccines known to those of skill in the art. The MTS can facilitate entry of the viral glycoprotein into the cell, where antigen processing of the protein can occur. Processed antigen expressed on the cell surface will provide an immune
30 response as has been seen in plasmid-delivery DNA vaccines, without the inherent limitations of plasmid-delivery systems.

An antigen-specific cytotoxic T lymphocyte (CTL) response has been demonstrated by the Naval Medical Research Institute for the circumsporozoite protein of *Plasmodium falciparum* (PfCSP) administered by means of naked plasmid DNA (Wang *et al.*, 1998). The MTS of the present invention and the method of engineering proteins for cell membrane permeability can be used to deliver purified fusion protein comprising the MTS and the PfCSP protein to provide antigenic stimulation of a CTL response. If provided in a suitable carrier, the MTS/PfCSP protein can be produced for oral or aerosol delivery in a more stable form which is appropriate for delivery and administration to subjects at remote locations, without costly and often unavailable measures usually associated with vaccine preservation.

Oral delivery of vaccines using the MTS fusion protein provides a method for delivering antigenic epitopes to the intestinal mucosa, where a strong CTL response can be generated. Alternatively, vaccines can be delivered using deep-lung delivery methods recently developed (Patton, 1997). Most proteins and peptides are absorbed naturally in the lungs, where they pass into the bloodstream. Protein absorption apparently occurs in the alveoli, by a process known as transcytosis. Use of deep-lung delivery methods would provide a non-invasive method of vaccine delivery for fusion proteins produced by the method of the present invention.

Genetically Engineered Proteins With Cell Membrane Permeability for Use in Drug Delivery Systems

New protein therapies have been developed for the treatment of previously untreatable conditions, including hepatitis C, hormonal disorders, multiple sclerosis, and some forms of cancer. Protein delivery into the extracellular

environment has presented a challenge, because of the large size and fragile three-dimensional structure of proteins. Protein stabilization has been accomplished by techniques as described, for example, in U.S. Patent No. 5,711,968, (incorporated herein by
5 reference) which describes the use of zinc to stabilize recombinant human growth hormone and recombinant -interferon in microspheres. U.S. Patent No. 5,674,534, issued to Zale, *et al.* (incorporated herein by reference), describes the use of ammonium sulfate to stabilize erythropoietin during release from hydrated
10 microspheres.

Methods of accomplishing sustained delivery of therapeutic protein products have also been described in U.S. Patent No. 4,767,628, issued to Hutchinson (incorporated herein by reference), and U.S. Patent No. 4,765,189, issued to Kent, *et al.*
15 (incorporated herein by reference). Controlled release microspheres, described in U.S. Patent No. 5,019,400, issued to Gombotz, *et al.* (incorporated herein by reference), and marketed as the ProLease[®] system (Alkermes, Inc.), provide a powder form of solid protein, homogeneously and rigidly dispersed within
20 porous polymer particles (often made of poly (lactide-co-glycolide), or PLG). An implantable osmotic pump system has also been reported to deliver peptide drugs at a constant rate for up to 1 year (Wright *et al.*, 1997).

The method of genetically engineering proteins with cell
25 membrane permeability described by the present invention provides a means for delivering therapeutic protein products into a cell. Combination of the present invention with previously described methods of extracellular protein delivery provide a method of delivering proteins for import into a cell in a stabilized,
30 functional form in a controlled-release fashion.

Polypeptides are produced using an appropriate expression vector and expression system. Cell membrane permeability is

conferred upon the protein or polypeptide by the expression of a fusion protein with the membrane translocating sequence (MTS) located either N-terminal or C-terminal to the expressed polypeptide. Less stable proteins are stabilized by methods known to those of skill in the art and described previously. Delivery to the extracellular environment is accomplished by providing the stabilized fusion protein in an appropriate carrier, such as the microsphere carriers described in U.S. Patent No. 5,019,400. The protein of choice will dictate the appropriate vector and expression system, as well as the appropriate stabilization and delivery technique. A person of skill in the art of drug delivery systems can choose the appropriate techniques from among those described.

Viral proteins which interfere with antigen presentation by down-regulating MHC class I expression have been identified for viruses such as herpesviruses, adenoviruses, and human immunodeficiency virus (Hengel *et al.* 1997). Use of these viral proteins to stabilize tissue transplants, such as beta cell transplants for diabetes treatment, against immune attack has been investigated. Gene delivery has been attempted through the use of viral vectors, however, with some success—but with the problems common to viral vector delivery systems. Viral vectors, introducing more protein material than the target protein, carry with them a risk of immune reaction to the vector itself. Particularly where more than one target protein must be delivered to more than one target tissue, these vectors can produce hypersensitivity reactions after repeated delivery. Using the method of the present invention, however, these disadvantages can be overcome. In the case of pancreatic beta cell transplant for diabetes, for example, a time-release drug delivery system can be provided using stabilizing agents such as those described in the references previously mentioned. The target protein for the drug delivery system can be a viral protein that inhibits MHC class I

expression on the transplanted cells. This protein can be produced as a fusion protein attached either N-terminal or C-terminal to an MTS as in the present invention. The MTS facilitates delivery of the viral protein to the interior of the cell, where the viral protein
5 inhibits MHC class I expression. In a preferred embodiment of the invention, the viral protein may be the E3 19K protein of human adenovirus.

During et al. demonstrated that lactose intolerance could be treated using peroral application of adeno-associated virus (AAV)
10 encoding the enzyme

Cancer Therapy Using Cell Permeable Proteins

The method of the present invention provides a means for producing cell-permeable proteins for the treatment of cancer.
15 Regulators of apoptosis and cell cycle control have been found to play a key role in oncogenesis, and gene therapy techniques using intratumoral injection of an adenoviral expression vector encoding the p53 gene have shown promise for the control of some tumors. Delivery of specific protein products through the use of viral
20 vectors has proven to be problematic, however. The MTS and method of the present invention provide a means for producing cell-permeable proteins from among the cell cycle regulators and regulators of apoptosis, as well as other proteins identified to play a role in the development of the cancer state.

For example, in the method of the present invention, the
25 nucleotide sequence encoding the p53 gene can be inserted into a suitable vector, either 5' or 3' to the sequence of an MTS of the present invention. Under expression conditions appropriate for the vector of choice and known to those of skill in the art, a fusion
30 protein comprising an MTS and the p53 protein can be expressed. Attachment of the MTS to the p53 protein renders the p53 protein cell permeable, and protein can be administered to tumor cells to

inhibit tumor development. Administration of cell-permeable protein can be accomplished in various ways, including, but not limited to, intratumoral injection, infusion, and intravenous administration. Bax and Bcl-xL are other examples from among a wide variety of proteins that have been determined to effect cell cycle control and apoptosis, and therefore be effective for cancer therapy. The method of the present invention provides a more efficient, less labor-intensive, less costly method for delivery of anti-oncogenic proteins to tumor cells.

Veterinary Applications of Proteins With Cell Membrane Permeability

A number of canine and feline diseases, as well as bovine and other diseases, provide attractive candidates for protein-based vaccines. Protein-based treatments for cancer and other disorders which have been developed for use in humans also provide therapeutic benefit in veterinary practice. A synthetic peptide vaccine for canine parvovirus (using the amino-terminal region of viral protein VP2), for example, has already proven to protect dogs against subsequent challenge with virulent canine parvovirus (Langeveld *et al.*, 1994).

Proteins With Cell Membrane Permeability Provide a More Effective Method for Vaccine Delivery Using Edible Plants as Vectors

Mason, *et al.* (1998) have described a method of using transgenic plants to provide an edible vaccine against enterotoxigenic *Escherichia coli* by expression of the heat-labile enterotoxin B subunit (LT-B). Agracetus has reported the development of transgenic corn, soybeans, and tobacco for production of recombinant proteins. In the method of the present

invention, corn, for example, can be genetically engineered, using recently developed techniques, to produce the fusion protein product of the MTS and the *E. coli* toxin LT-B. Transgenic corn fed to cattle will deliver the fusion protein to the lumen of the intestine, where the MTS will deliver the LT-B antigen to the interior of antigen-processing cells in the intestinal mucosa.

As discussed previously, antigenic proteins have been identified for a number of infectious agents. Methods for producing transgenic plants are known to those of skill in the relevant art, and have been described in the relevant literature. Expression of a synthetic fusion protein in potatoes, for example, under the control of a constitutive promoter provides an accumulation of the fusion protein in the tubers—and a supply of protein antigen for oral administration. Other edible plant species accumulate recombinant protein in leaves, which can be administered in a similar fashion to that previously described for the tubers of raw potatoes.

Antigen concentrations per gram of plant material are readily ascertainable by methods known to those of skill in the relevant art, which include protein isolation by standard means and quantification by standard protein assays. Oral antigen administration of microgram quantities of *E. coli* LT-B antigen, using genetically engineered raw potatoes, has been demonstrated to provide comparable protection in mice to antigen administration by conventional means.

Examples

Expression of the MTS fusion protein

In the present example, glutathione S-transferase from *Schistosoma japonicum* was used as the full-length protein cargo. Two different MTS expression plasmids were constructed, both utilizing the pGEX-3X vector (AMRAD Corporation Limited)

which encodes Sj26, the 26-kDa *S. japonicum* glutathione S-transferase (GST). Expression plasmid pGEX-3X-MTS1 and expression plasmid pGEX-3X-MTS2 were constructed so that the MTS coding sequence was located C-terminal to the Sj26 coding region (Fig. 3). Both plasmids contain a factor X cleavage site between GST and the MTS, which allows characterization of the attached MTS following enzyme cleavage. A *Bam*HI restriction endonuclease site located C-terminal to the MTS in pGEX-3X-MTS1 and N-terminal to the MTS in pGEX-3X-MTS2 provides an insertion point for a nucleotide sequence encoding a target polypeptide, protein domain, or full-length protein.

Wild-type GST (GST-WT), GST-MTS1, and GST-MTS2 were expressed in *E.coli* strain DH5 and purified from bacterial cell lysates by glutathione-agarose affinity chromatography. Analysis of the purified GST-MTS1 and GST-MTS2 protein preparations by SDS-PAGE showed predominantly single protein bands (29-kDa) with the predicted increase in apparent molecular size relative to GST-WT (Fig. 4). To confirm the amino acid content of the MTS in GST-MTS proteins, the GST-MTS1 protein was cleaved with factor Xa and the MTS-containing peptide was purified by high performance liquid chromatography (HPLC). The molecular weight (MW) of the purified MTS-containing peptide was determined by mass spectrometry analysis. The molecular weight of the purified MTS peptide matched the predicted value.

Cell Membrane Permeability of the MTS-Protein Fusion Product

Confluent NIH 3T3 cells were incubated with 20 M of purified GST-MTS1, with GST-WT and untreated cells used as control, in Dulbecco's Modified Eagles Medium (DMEM) for 30 minutes at 37°C. The cells were extensively washed and subsequently analyzed by indirect immunofluorescence assay

using anti-GST antibodies. In GST-MTS1 protein-treated cells, strong fluorescent signals were observed in a punctate staining pattern throughout all cells examined (Fig. 5a). GST-WT-treated cells did not display fluorescent signals (Fig. 5b). Other cell types
5 were also utilized, including murine endothelial LE-II cells and baby hamster kidney (BHK-21) cells, which exhibited a staining pattern similar to NIH 3T3 cells by the immunofluorescence assay.

Intracellular GST-MTS1 protein was also detected by Western blot analysis of cell lysates using polyclonal anti-GST
10 antibodies. A 29-kDa protein was detected by anti-GST antibodies in cell lysates from cells treated with GST-MTS1 (Fig. 6). GST was not detected in lysates from cells treated with GST-WT.

Flanking sequences of the MTS region did not affect the import activity conferred by the MTS, as GST-MTS2 cellular
15 import was as efficient as that of GST-MTS1.

GST-MTS1 protein was localized intracellularly, as evidenced by confocal laser scanning microscopy. A six-step Z-position 1 μ m sectional scanning of the cell exhibited strongest
20 fluorescent signals representing immunoreactive GST-MTS1 protein in the midsections of the cells (Fig. 7).

Influence of Protein Content, Time and Temperature of Incubation on Cellular Import

Western blot analysis was used to determine the relative
25 levels of protein imported into the cells. The import of GST-MTS1 protein by NIH3T3 cells at 37°C for 30 minutes incubation exhibited dose dependence over a concentration range up to 20 μ M (Fig. 8a). Higher concentrations have not yet been tested.

Uptake of GST-MTS1 by NIH 3T3 cells occurred equally
30 well at 22°C and 37°C (Fig. 8b). Protein import was significantly impaired at 4°C, however. At physiological temperatures, imported protein was readily detected after only 30 minutes

incubation. Protein import had not reached saturation after 6 hours of incubation. Stability of intracellular GST-MTS1 protein was evidence by the lack of degradation products after 6 hours incubation.

5 Imported protein was not localized within cellular compartments such as lysosomes, as evidenced by the observation that GST-MTS1 continued to accumulate intracellularly up to 18 hours incubation with only a low level of degraded protein detected (Fig. 8c). Furthermore, intact GST-MTS1 was recovered from
10 lysates of protein-treated cells using glutathione-conjugated agarose beads.

 Cells treated with GST-MTS1 at a final extracellular concentration of 20 M, for 30 minute incubation at 37°C, imported approximately $0.5-1 \times 10^6$ molecules of GST-MTS1
15 protein per cell, as determined by comparison of band intensity in Western blots of the protein recovered from cell lysates to that of known concentrations of the protein.

Lack of Cellular Cytotoxicity of the Protein Import

20 Protein-treated and untreated cells were stained with fluorescein diacetate-ethidium bromide. After 30 minutes incubation, the percentage of viable cells treated with 20 mM protein was 98.43 ± 0.96 , and after 2 hours the percentage of viable protein-treated cells was 98.40 ± 0.95 . In protein-untreated cells,
25 viability was 99.43 ± 0.39 and 97.90 ± 0.67 , respectively.

Use of Cellular Protein Import to Investigate Intracellular Signaling Processes

 A cell-permeable fusion protein containing the Grb2 SH2
30 domain (which binds to the tyrosine-phosphorylated EGF receptor) was constructed by synthesizing a DNA fragment encoding the human Grb2 SH2 domain (residues 54-164) using the polymerase

chain reaction (PCR) and inserting the sequence into GST-MTS2 and pGEX-3X (which contains the glutathione S-transferase coding sequence without the MTS). Protein products from each plasmid construct were expressed in *E. coli*, and purified. The MTS fusion protein was efficiently transported into SAA cells, as determined by indirect immunofluorescence spectroscopy (Fig. 9a and Fig. 9b) and Western blot analysis.

Serum-starved SAA cells (NIH 3T3 cells overexpressing epidermal growth factor receptor) were incubated with GST-Grb2SH2-MTS protein or control proteins for 1 hour prior to epidermal growth factor stimulation. Phosphorylated EGFR associated with endogenous Grb2 was examined in coimmunoprecipitation assays. Inducible EGFR/Grb2 association was inhibited in cells pretreated with GST-Grb2SH2-MTS (Fig. 10a). No significant inhibition was observed in cells pretreated with the non-cell permeable GST-Grb2SH2 protein or nonfunctional GST-MTS2 protein. Inhibition of EGFR/Grb2 association by GST-Grb2SH2-MTS was dose-dependent (Fig. 10b), reaching 35% at 0.4 M, 62% at 1 , and 93% at 2.5 final extracellular concentration as determined by densitometric analysis.

Pretreatment of cells with GST-Grb2SH2-MTS protein substantially inhibited the EGFR-induced MAP kinase activation involved in downstream mitogenic signaling. Specificity of binding of EGFR to GST-Grb2SH2-MTS was confirmed by coimmunoprecipitation assay with anti-GST antibodies.

The following references, to the extent that they provide details supplementary to those set forth herein, are specifically incorporated herein by reference:

Chen, *et al.* (1998), "Protective Immunity Induced by Oral Immunization with a Rotavirus DNA Vaccine Encapsulated in Microparticles," *J. Virol.* 72(7): 5757-5761.

- Chinsangaram, *et al.* (1998), "Antibody Response in Mice Inoculated with DNA Expressing Foot-and-Mouth Disease Virus Capsid Proteins," *J. Virol.* 72(5): 4454-4457.
- Derossi *et al.* (1994), "The Third Helix of the Antennapedia Homeodomain Translocates Through Biological Membranes," *J. Biol. Chem.* 269(14): 10444-10450.
- Dokka, *et al.* (1997), "Cellular Delivery of Oligonucleotides by Synthetic Import Peptide Carrier," *Pharmaceutical Research* 14: 1759-1764.
- Griffin, *et al.* (1998), "Specific Covalent Labeling of Recombinant Protein Molecules Inside Living Cells," *Science* 281: 269-272.
- Hengel, H. *et al.* (1997), "Interference with Antigen Processing by Viruses," *Curr. Opin. Immunol.* 9(4): 470-6.
- Jones, *et al.* (1998), "Role of Dynamin in the Formation of Transport Vesicles from the Trans-Golgi Network," *Science* 279: 573-577.
- King, *et al.* (1981), "Biochemical Identification of Viruses Causing the 1981 Outbreaks of Foot-and-Mouth Disease in the UK," *Nature* 293: 479-480.
- Kohler, *et al.* (1997), "Exchange of Protein Molecules Through Connections Between Higher Plant Plastids," *Science* 276: 2039-2042.
- Langeveld, *et al.* (1994), "First Peptide Vaccine Providing Protection Against Viral Infection in the Target Animal: Studies of Canine Parvovirus in Dogs," *J. Virol.* 68(7): 4506-4513.
- Lewin, Benjamin (1994), *Genes V*, Oxford University Press, p. 290.
- Lin *et al.* (1995), *J. Biol. Chem.* 270(24): 14255-14258.
- Mason, *et al.* (1998), "Edible Vaccine Protects Mice Against *Escherichia coli* Heat-Labile Enterotoxin (LT): Potatoes Expressing a Synthetic LT-B Gene," *Vaccine* 16(13): 1336-1343.
- Mathews, *et al.* (1992), "A Synthetic Peptide to the E Glycoprotein of Murray Valley Encephalitis Virus Defines Multiple Virus-

- Reactive T- and B-cell Epitopes," *J. Virol.* 66(11): 6555-6562.
- No, *et al.* (1996), "Ecdysone-inducible Gene Expression in Mammalian Cells and Transgenic Mice," *Proc. Natl. Acad. Sci. USA* 93: 3346-3351.
- 5 Olson, *et al.* (1995), "Analysis of MAP4 Function in Living Cells Using Green Fluorescent Protein (GFP) Chimeras," *J. Cell. Biol.* 130(3): 639-50.
- Patton, J. (1997), "Deep-lung Delivery of Therapeutic Proteins," *CHEMTECH* 27(12): 34-38.
- 10 Tam, *et al.* (1995), "Peptide Synthesis Using Unprotected Peptides Through Orthogonal Coupling Methods," *Proc. Natl. Acad. Sci. USA* 92: 12485-12489.
- Tsien, R.Y. (1998), *Annu. Rev. Biochem.* 67: 509.
- Wang, *et al.* (1998), "Induction of Antigen-Specific Cytotoxic T
- 15 Lymphocytes in Humans by a Malaria DNA Vaccine," *Science* 282: 476-480.
- Wright, *et al.* (1997), *Proc. Int. Symp. Controlled Release Bioact. Mater.* 24: 59.
- Xu, *et al.* (1998), "Immunization for Ebola Virus Infection," *Nat. Med.* 4(1): 37-42.
- 20

SEQUENCE LISTING

(1) GENERAL INFORMATION

(i) APPLICANT: Lin, Yao-Zhong
Donahue, John P.
5 Rojas, Mauricio
Tan, Zhongjia

(ii) TITLE OF INVENTION: "Sequence and Method for
Genetic Engineering of Proteins with Cell Membrane
Translocating Activity"

10 (iii) NUMBER OF SEQUENCES: 18

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Wadley & Patterson
(B) STREET: Suite 2020 NationsBank
Plaza
15 414 Union Street
(C) CITY: Nashville
(D) STATE: Tennessee
(E) COUNTRY: USA
(F) ZIP CODE: 37219

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: 1.44MB High
Density Diskette
(B) COMPUTER TYPE: IBM Compatible
25 (C) OPERATING SYSTEM: Windows 98
(D) SOFTWARE: PatIn Version
2.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: Not yet known
30 (B) FILING DATE: 4 November 1998
(C) CLASSIFICATION:

- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: US 60/080,083
- (B) FILING DATE: 31 March 1998
- (viii) ATTORNEY/AGENT INFORMATION:
- 5 (A) NAME: Patterson, Mark J.
- (B) REGISTRATION NUMBER: 30,412
- (C) REFERENCE/DOCKET NUMBER: 3219
- (ix) TELECOMMUNICATION INFORMATION:
- 10 (A) TELEPHONE: (615) 242-2400
- (B) TELEFAX: (615) 242-2221
- (2) INFORMATION FOR SEQ ID NO:1
- (i) SEQUENCE CHARACTERISTICS:
- 15 (A) LENGTH: 12 amino acid residues
- (B) TYPE: amino acid
- (C) STRANDEDNESS: Not applicable
- (D) TOPOLOGY: Not applicable
- (ii) MOLECULE TYPE: Peptide
- 20 (iii) HYPOTHETICAL: No
- (iv) ANTI-SENSE: No
- (v) FRAGMENT TYPE: Not applicable (artificial
sequence)
- (vi) ORIGINAL SOURCE: Artificial (synthesized)
- 25 (vii) IMMEDIATE SOURCE: Artificial (synthesized)
- (viii) POSITION IN GENOME: Not applicable
- (ix) FEATURE: Peptide translocates proteins across
cell membrane
- (A) NAME/KEY: Membrane translocation peptide
- 30 sequence
- (B) LOCATION: Amino acid residues 1-12

(C) IDENTIFICATION METHOD: Experimental

(D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:

5
Rojas, Mauricio
Donahue, John
Tan, Zhonjia
Lin, Yao-Zhong

10
(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

(D) VOLUME: 16

(E) ISSUE: April 1998

(F) PAGES: 370-375

15
(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Ala Ala Val Leu Leu Pro Val Leu Leu Ala Ala Pro

20 (3) INFORMATION FOR SEQ ID NO: 2

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 8 amino acid residues

(B) TYPE: amino acid

(C) STRANDEDNESS: Not applicable

25 (D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: Peptide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

30 (v) FRAGMENT TYPE: Not applicable (artificial
sequence)

(vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

(viii) POSITION IN GENOME: Not applicable

(ix) FEATURE: Peptide translocates proteins across
cell membrane

(A) NAME/KEY: Membrane translocation peptide

5 sequence

(B) LOCATION: Amino acid residues 1-8

(C) IDENTIFICATION METHOD: Experimental

(D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

10

(A) AUTHORS:

Rojas, Mauricio

Donahue, John

Tan, Zhongjia

Lin, Yao-Zhong

15

(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

(D) VOLUME: 16

(E) ISSUE: April 1998

20

(F) PAGES: 370-375

(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Ala Ala Val Leu Leu Pro Val Leu

25

(4) INFORMATION FOR SEQ ID NO: 3

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 amino acid residues

(B) TYPE: amino acid

(C) STRANDEDNESS: Not applicable

30

(D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: Peptide

(iii) HYPOTHETICAL: No

- (iv) ANTI-SENSE: No
- (v) FRAGMENT TYPE: Not applicable (artificial
sequence)
- (vi) ORIGINAL SOURCE: Artificial (synthesized)
- 5 (vii) IMMEDIATE SOURCE: Artificial (synthesized)
- (viii) POSITION IN GENOME: Not applicable
- (ix) FEATURE: Peptide translocates proteins across
cell membrane
- 10 (A) NAME/KEY: Membrane translocation peptide
sequence
- (B) LOCATION: Amino acid residues 1-9
- (C) IDENTIFICATION METHOD: Experimental
- (D) OTHER INFORMATION:
- 15 (x) PUBLICATION INFORMATION:
- (A) AUTHORS:
- Rojas, Mauricio
- Donahue, John
- 20 Tan, Zhonjia
- Lin, Yao-Zhong
- (B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"
- 25 (C) JOURNAL: Nature Biotechnology
- (D) VOLUME: 16
- (E) ISSUE: April 1998
- (F) PAGES: 370-375
- (G) DATE: 01-APR-1998
- 30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:
Ala Ala Val Leu Leu Pro Val Leu Leu

(E) ISSUE: April 1998
(F) PAGES: 370-375
(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

5 Ala Ala Val Leu Leu Pro Val Leu Leu Ala

(6) INFORMATION FOR SEQ ID NO: 5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 amino acid residues
(B) TYPE: amino acid
10 (C) STRANDEDNESS: Not applicable
(D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: Peptide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

15 (v) FRAGMENT TYPE: Not applicable (artificial
sequence)

(vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

(viii) POSITION IN GENOME: Not applicable

20 (ix) FEATURE: Peptide translocates proteins across
cell membrane

(A) NAME/KEY: Membrane translocation peptide
sequence

25 (B) LOCATION: Amino acid residues 1-11

(C) IDENTIFICATION METHOD: Experimental

(D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:

30 Rojas, Mauricio
Donahue, John
Tan, Zhonjia

(5) INFORMATION FOR SEQ ID NO: 4

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acid residues

(B) TYPE: amino acid

5 (C) STRANDEDNESS: Not applicable

(D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: Peptide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

10 (v) FRAGMENT TYPE: Not applicable (artificial
sequence)

(vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

(viii) POSITION IN GENOME: Not applicable

15 (ix) FEATURE: Peptide translocates proteins across
cell membrane

(A) NAME/KEY: Membrane translocation peptide

sequence

20 (B) LOCATION: Amino acid residues 1-10

(C) IDENTIFICATION METHOD: Experimental

(D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:

25 Rojas, Mauricio

Donahue, John

Tan, Zhonjia

Lin, Yao-Zhong

30 (B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

(D) VOLUME: 16

Lin, Yao-Zhong

(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

5 (D) VOLUME: 16

(E) ISSUE: April 1998

(F) PAGES: 370-375

(G) DATE: 01-APR-1998

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Ala Ala Val Leu Leu Pro Val Leu Leu Ala Ala

(7) INFORMATION FOR SEQ ID NO: 6

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 8 amino acid residues

(B) TYPE: amino acid

(C) STRANDEDNESS: Not applicable

(D) TOPOLOGY: Not applicable

20 (ii) MOLECULE TYPE: Peptide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(v) FRAGMENT TYPE: Not applicable (artificial
sequence)

25 (vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

(viii) POSITION IN GENOME: Not applicable

(ix) FEATURE: Peptide translocates proteins across
cell membrane

30 (A) NAME/KEY: Membrane translocation peptide
sequence

(B) LOCATION: Amino acid residues 1-8

(C) IDENTIFICATION METHOD: Experimental

(D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:

5
Rojas, Mauricio
Donahue, John
Tan, Zhonjia
Lin, Yao-Zhong

10
(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

(D) VOLUME: 16

(E) ISSUE: April 1998

15
(F) PAGES: 370-375

(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Leu Pro Val Leu Leu Ala Ala Pro

(8) INFORMATION FOR SEQ ID NO: 7

20
(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 amino acid residues

(B) TYPE: amino acid

(C) STRANDEDNESS: Not applicable

(D) TOPOLOGY: Not applicable

25
(ii) MOLECULE TYPE: Peptide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(v) FRAGMENT TYPE: Not applicable (artificial
sequence)

30
(vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

(viii) POSITION IN GENOME: Not applicable

(ix) FEATURE: Peptide translocates proteins across
cell membrane

(A) NAME/KEY: Membrane translocation peptide
sequence

5 (B) LOCATION: Amino acid residues 1-9
(C) IDENTIFICATION METHOD: Experimental
(D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:
10 Rojas, Mauricio
Donahue, John
Tan, Zhonjia
Lin, Yao-Zhong

15 (B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology
(D) VOLUME: 16
(E) ISSUE: April 1998
20 (F) PAGES: 370-375
(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:
Leu Leu Pro Val Leu Leu Ala Ala Pro

25 (9) INFORMATION FOR SEQ ID NO: 8

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acid residues
(B) TYPE: amino acid
(C) STRANDEDNESS: Not applicable
30 (D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: Peptide

(iii) HYPOTHETICAL: No

- (iv) ANTI-SENSE: No
- (v) FRAGMENT TYPE: Not applicable (artificial sequence)
- (vi) ORIGINAL SOURCE: Artificial (synthesized)
- 5 (vii) IMMEDIATE SOURCE: Artificial (synthesized)
- (viii) POSITION IN GENOME: Not applicable
- (ix) FEATURE: Peptide translocates proteins across cell membrane
- (A) NAME/KEY: Membrane translocation peptide
- 10 sequence
- (B) LOCATION: Amino acids 1-10
- (C) IDENTIFICATION METHOD: Experimental
- (D) OTHER INFORMATION:
- (x) PUBLICATION INFORMATION:
- 15 (A) AUTHORS:
- Rojas, Mauricio
- Donahue, John
- Tan, Zhonjia
- Lin, Yao-Zhong
- 20
- (B) TITLE: "Genetic Engineering of Proteins with Cell Membrane Permeability"
- (C) JOURNAL: Nature Biotechnology
- (D) VOLUME: 16
- 25 (E) ISSUE: April 1998
- (F) PAGES: 370-375
- (G) DATE: 01-APR-1998
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:
- Val Leu Leu Pro Val Leu Leu Ala Ala Pro
- 30 (10) INFORMATION FOR SEQ ID NO: 9
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 11 amino acid residues

(B) TYPE: amino acid
(C) STRANDEDNESS: Not applicable
(D) TOPOLOGY: Not applicable
(ii) MOLECULE TYPE: Peptide
5 (iii) HYPOTHETICAL: No
(iv) ANTI-SENSE: No
(v) FRAGMENT TYPE: Not applicable (artificial
sequence)
(vi) ORIGINAL SOURCE: Artificial (synthesized)
10 (vii) IMMEDIATE SOURCE: Artificial (synthesized)
(viii) POSITION IN GENOME: Not applicable
(ix) FEATURE: Peptide translocates proteins across
cell membrane
(A) NAME/KEY: Membrane translocation peptide
15 sequence
(B) LOCATION: Amino acid residues 1-11
(C) IDENTIFICATION METHOD: Experimental
(D) OTHER INFORMATION:
(x) PUBLICATION INFORMATION:
20 (A) AUTHORS:
Rojas, Mauricio
Donahue, John
Tan, Zhonjia
Lin, Yao-Zhong
25
(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"
(C) JOURNAL: Nature Biotechnology
(D) VOLUME: 16
30 (E) ISSUE: April 1998
(F) PAGES: 370-375
(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Ala Val Leu Leu Pro Val Leu Leu Ala Ala Pro

5 (11) INFORMATION FOR SEQ ID NO: 10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 nucleotide residues

(B) TYPE: deoxyribonucleic acid

(C) STRANDEDNESS: Double-stranded artificial

10 sequence

(D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: polynucleotide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

15 (v) FRAGMENT TYPE: Not applicable (artificial
sequence)

(vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

(viii) POSITION IN GENOME: Not applicable

20 (ix) FEATURE: Nucleotide sequence of peptide that
translocates proteins across

cell membrane

(A) NAME/KEY: Membrane translocation peptide
polynucleotide coding sequence

25 (B) LOCATION: Nucleotide residues 1-36

(C) IDENTIFICATION METHOD: Experimental

(D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:

30 Rojas, Mauricio
Donahue, John
Tan, Zhonjia

Lin, Yao-Zhong

(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

5 (C) JOURNAL: Nature Biotechnology
(D) VOLUME: 16
(E) ISSUE: April 1998
(F) PAGES: 370-375
(G) DATE: 01-APR-1998

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

gcagccgttc ttctccctgt tcttcttgcc gcaccc

15 (12) INFORMATION FOR SEQ ID NO: 11

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 nucleotide residues
(B) TYPE: deoxyribonucleic acid
(C) STRANDEDNESS: Double-stranded artificial

20 sequence

(D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: polynucleotide

(iii) HYPOTHETICAL: No

25 (iv) ANTI-SENSE: No

(v) FRAGMENT TYPE: Not applicable (artificial
sequence)

(vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

30 (viii) POSITION IN GENOME: Not applicable

(ix) FEATURE: Nucleotide sequence of peptide that
translocates proteins across

cell membrane

(A) NAME/KEY: Membrane translocation peptide
polynucleotide coding sequence

5

(B) LOCATION: Nucleotide residues 1-24

(C) IDENTIFICATION METHOD: Experimental

(D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:

10

Rojas, Mauricio

Donahue, John

Tan, Zhonjia

Lin, Yao-Zhong

15

(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

(D) VOLUME: 16

(E) ISSUE: April 1998

20

(F) PAGES: 370-375

(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

gcagccgttc ttctccctgt tctt

(13) INFORMATION FOR SEQ ID NO: 12

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 nucleotide residues

(B) TYPE: deoxyribonucleic acid

(C) STRANDEDNESS: Double-stranded artificial

sequence

30

(D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: polynucleotide

- (iii) HYPOTHETICAL: No
- (iv) ANTI-SENSE: No
- (v) FRAGMENT TYPE: Not applicable (artificial
5 sequence)
- (vi) ORIGINAL SOURCE: Artificial (synthesized)
- (vii) IMMEDIATE SOURCE: Artificial (synthesized)
- (viii) POSITION IN GENOME: Not applicable
- (ix) FEATURE: Nucleotide sequence of peptide that
10 translocates proteins across
cell membrane
- (A) NAME/KEY: Membrane translocation peptide
polynucleotide coding sequence
- 15 (B) LOCATION: Nucleotide residues 1-27
- (C) IDENTIFICATION METHOD: Experimental
- (D) OTHER INFORMATION:
- (x) PUBLICATION INFORMATION:
- (A) AUTHORS:
- 20 Rojas, Mauricio
Donahue, John
Tan, Zhonjia
Lin, Yao-Zhong
- 25 (B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"
- (C) JOURNAL: Nature Biotechnology
- (D) VOLUME: 16
- (E) ISSUE: April 1998
- 30 (F) PAGES: 370-375
- (G) DATE: 01-APR-1998
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

gcagccgttc ttctccctgt tcttctt

(14) INFORMATION FOR SEQ ID NO: 13

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 30 nucleotide residues
(B) TYPE: deoxyribonucleic acid
(C) STRANDEDNESS: Double-stranded artificial

sequence

(D) TOPOLOGY: Not applicable

- 10 (ii) MOLECULE TYPE: polynucleotide
(iii) HYPOTHETICAL: No
(iv) ANTI-SENSE: No
(v) FRAGMENT TYPE: Not applicable (artificial
sequence)

- 15 (vi) ORIGINAL SOURCE: Artificial (synthesized)
(vii) IMMEDIATE SOURCE: Artificial (synthesized)
(viii) POSITION IN GENOME: Not applicable
(ix) FEATURE: Nucleotide sequence of peptide that
translocates proteins across

20 cell membrane

(A) NAME/KEY: Membrane translocation peptide
polynucleotide coding sequence

(B) LOCATION: Nucleotide residues 1-30

(C) IDENTIFICATION METHOD: Experimental

25 (D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:

Rojas, Mauricio

Donahue, John

30 Tan, Zhonjia

Lin, Yao-Zhong

(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

(D) VOLUME: 16

5 (E) ISSUE: April 1998

(F) PAGES: 370-375

(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

10

gcagccgttc ttctccctgt tcttcttgcc

(15) INFORMATION FOR SEQ ID NO: 14

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 nucleotide residues

(B) TYPE: deoxyribonucleic acid

(C) STRANDEDNESS: Double-stranded artificial
sequence

20 (D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: polynucleotide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

25 (v) FRAGMENT TYPE: Not applicable (artificial
sequence)

(vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

(viii) POSITION IN GENOME: Not applicable

30 (ix) FEATURE: Nucleotide sequence of peptide that
translocates proteins across
cell membrane

(A) NAME/KEY: Membrane translocation peptide
polynucleotide coding sequence

5 (B) LOCATION: Nucleotide residues 1-33
(C) IDENTIFICATION METHOD: Experimental
(D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:
10 Rojas, Mauricio
Donahue, John
Tan, Zhonjia
Lin, Yao-Zhong

15 (B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology
(D) VOLUME: 16
(E) ISSUE: April 1998
20 (F) PAGES: 370-375
(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

25 gcagccgttc ttctccctgt tcttcttgcc gca

(16) INFORMATION FOR SEQ ID NO: 15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 nucleotide residues
30 (B) TYPE: deoxyribonucleic acid
(C) STRANDEDNESS: Double-stranded artificial
sequence

- (D) TOPOLOGY: Not applicable
- (ii) MOLECULE TYPE: polynucleotide
- (iii) HYPOTHETICAL: No
- (iv) ANTI-SENSE: No
- 5 (v) FRAGMENT TYPE: Not applicable (artificial sequence)
- (vi) ORIGINAL SOURCE: Artificial (synthesized)
- (vii) IMMEDIATE SOURCE: Artificial (synthesized)
- (viii) POSITION IN GENOME: Not applicable
- 10 (ix) FEATURE: Nucleotide sequence of peptide that translocates proteins across cell membrane
- (A) NAME/KEY: Membrane translocation peptide
- 15 polynucleotide coding sequence
- (B) LOCATION: Nucleotide residues 1-24
- (C) IDENTIFICATION METHOD: Experimental
- (D) OTHER INFORMATION:
- (x) PUBLICATION INFORMATION:
- 20 (A) AUTHORS:
- Rojas, Mauricio
- Donahue, John
- Tan, Zhonjia
- Lin, Yao-Zhong
- 25 (B) TITLE: "Genetic Engineering of Proteins with Cell Membrane Permeability"
- (C) JOURNAL: Nature Biotechnology
- (D) VOLUME: 16
- 30 (E) ISSUE: April 1998
- (F) PAGES: 370-375

(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

5 cccccctgttc ttctttgccgc accc

(17) INFORMATION FOR SEQ ID NO: 16

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 27 nucleotide residues
(B) TYPE: deoxyribonucleic acid
(C) STRANDEDNESS: Double-stranded artificial
sequence

(D) TOPOLOGY: Not applicable

15 (ii) MOLECULE TYPE: polynucleotide
(iii) HYPOTHETICAL: No
(iv) ANTI-SENSE: No
(v) FRAGMENT TYPE: Not applicable (artificial sequence)

20 (vi) ORIGINAL SOURCE: Artificial (synthesized)
 (vii) IMMEDIATE SOURCE: Artificial (synthesized)
 (viii) POSITION IN GENOME: Not applicable
 (ix) FEATURE: Nucleotide sequence of peptide that
 translocates proteins across
 25 cell membrane

(A) NAME/KEY: Membrane translocation peptide
polynucleotide coding sequence

(B) LOCATION: Nucleotide residues 1-27

(C) IDENTIFICATION METHOD: Experimental

30 (D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:

Rojas, Mauricio

Donahue, John

Tan, Zhonjia

Lin, Yao-Zhong

5

(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

(D) VOLUME: 16

10

(E) ISSUE: April 1998

(F) PAGES: 370-375

(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

15

ctctccctg ttctcttgc cgcaccc

(18) INFORMATION FOR SEQ ID NO: 17

(i) SEQUENCE CHARACTERISTICS:

20

(A) LENGTH: 30 nucleotide residues

(B) TYPE: deoxyribonucleic acid

(C) STRANDEDNESS: Double-stranded artificial

sequence

(D) TOPOLOGY: Not applicable

25

(ii) MOLECULE TYPE: polynucleotide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(v) FRAGMENT TYPE: Not applicable (artificial

30

sequence)

(vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

(viii) POSITION IN GENOME: Not applicable

(ix) FEATURE: Nucleotide sequence of peptide that
translocates proteins across
cell membrane

5 (A) NAME/KEY: Membrane translocation peptide
polynucleotide coding sequence

(B) LOCATION: Nucleotide residues 1-30

(C) IDENTIFICATION METHOD: Experimental

(D) OTHER INFORMATION:

10 (x) PUBLICATION INFORMATION:

(A) AUTHORS:

Rojas, Mauricio

Donahue, John

Tan, Zhonjia

15 Lin, Yao-Zhong

(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

(D) VOLUME: 16

20 (E) ISSUE: April 1998

(F) PAGES: 370-375

(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

gtttcttctcc ctgttcttct tgccgcaccc

25 (19) INFORMATION FOR SEQ ID NO: 18

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 nucleotide residues

(B) TYPE: deoxyribonucleic acid

(C) STRANDEDNESS: Double-stranded artificial

30 sequence

(D) TOPOLOGY: Not applicable

(ii) MOLECULE TYPE: polynucleotide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(v) FRAGMENT TYPE: Not applicable (artificial sequence)

5 (vi) ORIGINAL SOURCE: Artificial (synthesized)

(vii) IMMEDIATE SOURCE: Artificial (synthesized)

(viii) POSITION IN GENOME: Not applicable

(ix) FEATURE: Nucleotide sequence of peptide that
translocates proteins across

10 cell membrane

(A) NAME/KEY: Membrane translocation peptide
polynucleotide coding sequence

(B) LOCATION: Nucleotide residues 1-33

(C) IDENTIFICATION METHOD: Experimental

15 (D) OTHER INFORMATION:

(x) PUBLICATION INFORMATION:

(A) AUTHORS:

Rojas, Mauricio

Donahue, John

20 Tan, Zhonjia

Lin, Yao-Zhong

(B) TITLE: "Genetic Engineering of Proteins with
Cell Membrane Permeability"

(C) JOURNAL: Nature Biotechnology

25 (D) VOLUME: 16

(E) ISSUE: April 1998

(F) PAGES: 370-375

(G) DATE: 01-APR-1998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

30 gccgttcttc tccctgttct tcttgccgca ccc

CLAIMS

What is claimed is:

1. An isolated peptide of about 8 to about 50 residues comprising at least eight consecutive residues of SEQ. ID NO. 1:
5 Ala-Ala-Val-Leu-Leu-Pro-Val-Leu-Leu-Ala-Ala-Pro.
2. The isolated peptide of claim 1 further comprising at least nine consecutive residues of SEQ. ID NO. 1.
3. The isolated peptide of claim 1 further comprising at least ten consecutive residues of SEQ. ID NO. 1.
- 10 4. The isolated peptide of claim 1 further comprising at least eleven consecutive residues of SEQ. ID NO. 1.
5. The isolated peptide of claim 1 further comprising at least twelve consecutive residues of SEQ. ID NO. 1.
- 15 6. An isolated polynucleotide comprising a sequence encoding at least eight consecutive residues of SEQ. ID NO. 1.
7. The isolated polynucleotide of claim 6 wherein the sequence encodes at least nine consecutive residues of SEQ. ID NO. 1.
- 20 8. The isolated polynucleotide of claim 6 wherein the sequence encodes at least ten consecutive residues of SEQ. ID NO. 1.
9. The isolated polynucleotide of claim 6 wherein the sequence encodes at least eleven consecutive residues of SEQ. ID NO. 1.
- 25 10. The isolated polynucleotide of claim 6 wherein the sequence encodes at least twelve consecutive residues of SEQ. ID NO. 1.
11. An isolated fusion polypeptide comprising
 - a) a peptide of 8 to 50 residues comprising at
30 least eight consecutive residues of SEQ. ID NO. 1:
Ala-Ala-Val-Leu-Leu-Pro-Val-Leu-Leu-Ala-Ala-Pro,
and

b) a cargo polypeptide.

12. The isolated fusion polypeptide of claim 11 wherein the peptide of about 8 to about 50 residues comprises at least nine consecutive residues of SEQ. ID NO. 1.

5 13. The isolated fusion polypeptide of claim 11 wherein the peptide of about 8 to about 50 residues comprises at least ten consecutive residues of SEQ. ID NO. 1.

10 14. The isolated fusion polypeptide of claim 11 wherein the peptide of about 8 to about 50 residues comprises at least eleven consecutive residues of SEQ. ID NO. 1.

15 15. The isolated fusion polypeptide of claim 11 wherein the peptide of about 8 to about 50 residues comprises at least twelve consecutive residues of SEQ. ID NO. 1.

15 16. The isolated fusion polypeptide of claim 11 wherein the cargo polypeptide further comprises a peptide.

17. The isolated fusion polypeptide of claim 11 wherein the cargo polypeptide further comprises a polypeptide.

18. The isolated fusion polypeptide of claim 11 wherein the cargo polypeptide further comprises a protein domain.

20 19. The isolated fusion polypeptide of claim 11 wherein the cargo polypeptide further comprises a protein.

20. The isolated fusion polypeptide of claim 11 wherein the cargo polypeptide further comprises a complex formed by two or more polypeptides.

25 21. The isolated fusion polypeptide of claim 11 wherein the cargo polypeptide further comprises a cell cycle regulatory protein.

22. The isolated fusion polypeptide of claim 11 wherein the cargo polypeptide further comprises an intracellular enzyme.

30 23. An isolated expression construct comprising

 a) a polynucleotide including a sequence encoding at least eight consecutive residues of SEQ. ID NO. 1 and

b) a promoter positioned to direct the transcription of said polynucleotide.

24. The isolated expression construct of claim 23, further comprising a multipurpose cloning site 5' to the polynucleotide.

5 25. The isolated expression construct of claim 23, further comprising a multipurpose cloning site 3' to the polynucleotide.

26. The isolated expression construct of claim 23, wherein the promoter comprises the bacterial tac promoter.

10 27. The isolated expression construct of claim 23, wherein the promoter comprises the bacterial phage T7 promoter.

28. The isolated expression construct of claim 23 wherein the promoter comprises the cytomegalovirus promoter.

29. The isolated expression construct of claim 23 wherein the promoter comprises the respiratory syncytial virus promoter.

15 30. The isolated expression construct of claim 23 wherein the promoter comprises the *Pichia pastoris* AOX1 promoter.

31. The isolated expression construct of claim 23 wherein the promoter comprises an element of the pGEX DNA expression vector.

20 32. The isolated expression construct of claim 23, further comprising a polynucleotide sequence encoding an affinity purification tag.

25 33. The isolated expression construct of claim 32 wherein the affinity purification tag comprises six consecutive histidine amino acid residues.

34. The isolated expression construct of claim 32 wherein the affinity purification tag comprises the glutathione S-transferase domain.

30 35. The isolated expression construct of claim 32 further comprising a nucleotide sequence encoding a factor X site positioned between the affinity purification tag and the polynucleotide comprising a sequence encoding at least eight

consecutive residues of SEQ. ID NO. 1.

36. A method of producing a cell-permeable polypeptide comprising introducing into a suitable host cell an expression construct comprising:

- 5 a) a polynucleotide including a sequence encoding at least eight consecutive residues of SEQ. ID NO. 1 fused to a sequence encoding a polypeptide of interest, and
- b) a promoter positioned to direct the transcription of said polynucleotide.

10 37. The method of claim 36 wherein the polynucleotide further comprises a sequence encoding at least nine consecutive residues of SEQ. ID NO. 1.

 38. The method of claim 36 wherein the polynucleotide further comprises a sequence encoding at least ten consecutive
15 residues of SEQ. ID NO. 1.

 39. The method of claim 36 wherein the polynucleotide further comprises a sequence encoding at least eleven consecutive residues of SEQ. ID NO. 1.

 40. The method of claim 36 wherein the polynucleotide
20 further comprises a sequence encoding at least twelve consecutive residues of SEQ. ID NO. 1.

 41. The method of claim 36 wherein the polypeptide of interest comprises *Schistosoma japonicum* glutathione S-transferase.

25 42. The method of claim 36 wherein the polypeptide of interest comprises the adenovirus E3 19K protein.

 43. The method of claim 36 wherein the polypeptide of interest comprises the mammalian p53 protein.

 44. A method of increasing cellular permeability of a
30 molecule comprising attaching the molecule to a peptide of 8 to 50 amino acids including at least 8 amino acids of SEQ. ID NO. 1.

 45. The method of claim 44 wherein the peptide of 8 to 50

amino acids comprises at least 9 amino acids of SEQ. ID NO. 1.

46. The method of claim 44 wherein the peptide of 8 to 50 amino acids comprises at least 10 amino acids of SEQ. ID NO. 1.

47. The method of claim 44 wherein the peptide of 8 to 50 amino acids comprises at least 11 amino acids of SEQ. ID NO. 1.

48. The method of claim 44 wherein the peptide of 8 to 50 amino acids comprises at least 12 amino acids of SEQ. ID NO. 1.

49. The method of claim 44 wherein attaching the molecule to the peptide comprises attachment of the molecule N-terminal to the peptide.

50. The method of claim 44 wherein attaching the molecule to the peptide comprises attachment of the molecule C-terminal to the peptide.

51. The method of claim 44 wherein attaching the molecule to the peptide comprises attachment by a peptide bond.

52. The method of claim 44 wherein attaching the molecule to the peptide comprises attachment by a covalent bond.

53. The method of claim 52 wherein attachment by a covalent bond comprises attachment by a poly-L-lysine linker sequence.

54. The method of claim 52 wherein attachment by a covalent bond comprises attachment by orthogonal coupling.

55. A method for inducing an immune response in a mammalian subject comprising administering to the animal an immunogenic dose of a cell-permeable fusion protein including

- a) a peptide of 8 to 50 residues comprising at least eight consecutive residues of SEQ. ID NO. 1 and
- b) a polypeptide.

56. The method of claim 55 wherein the peptide comprises at least 9 consecutive residues of SEQ. ID NO. 1.

57. The method of claim 55 wherein the peptide comprises at least 10 consecutive residues of SEQ. ID NO. 1.

58. The method of claim 55 wherein the peptide comprises at least 11 consecutive residues of SEQ. ID NO. 1.

59. The method of claim 55 wherein the peptide comprises at least 12 consecutive residues of SEQ. ID NO. 1.

5 60. The method of claim 55 wherein the polypeptide comprises a viral polypeptide.

61. The method of claim 60 wherein the viral polypeptide comprises the hepatitis B surface antigen.

62. A method of protecting a subject from an infectious agent comprising administering to the subject a cell-permeable fusion protein including

a) a peptide of 8 to 50 residues comprising at least eight consecutive residues of SEQ. ID NO. 1 and

b) a polypeptide that inhibits reproduction of the infectious agent.

63. The method of claim 62 wherein the polypeptide comprises a protease inhibitor.

64. A method for treating cancer in a subject comprising administering to the subject a cell-permeable fusion protein including

a) a peptide of 8 to 50 residues comprising at least eight consecutive residues of SEQ. ID NO. 1 and

b) a polypeptide tumor suppressor.

65. The method of claim 64 wherein the polypeptide comprises a regulator of cell cycle progression.

66. The method of claim 65 wherein the polypeptide comprises the p53 protein.

67. The method of claim 63 wherein the polypeptide comprises an inhibitor of Bcl-2.

68. A method of genetically engineering labeled proteins for membrane permeability, comprising construction of a polynucleotide vector to express a chimeric protein including:

a) a peptide of 8 to 50 residues comprising at least eight consecutive residues of SEQ. ID NO. 1;

b) a molecular label; and

c) a target protein.

5 69. The method of claim 68 wherein the molecular label further comprises a green fluorescent protein of *Aequora victoria*.

70. The method of claim 68 wherein the molecular label further comprises a blue fluorescent protein.

10 71. The method of claim 68 wherein the target protein further comprises a MAP kinase.

72. The method of claim 68 wherein the target protein further comprises a caspase.

73. The method of claim 68 wherein the target protein further comprises Bcl-2.

15

1/9

	<u>Factor X</u>	<u>MTS</u>	<u>BamHI</u>	<u>SmaI</u>	<u>EcoRI</u>
MTS1	ATCGAAGTCTGGGATCGCAGCCGTTCTCTCCCTGTTCTTCCCGCACCCCGGGATCCCCGGGAATTCATCGTGA I E G R G I A A V L L P V L L A A P G I P G N S S ***				
	<u>Factor X</u>	<u>MTS</u>			<u>EcoRI</u>
MTS2	ATCGAAGTCTGGGATCCCCGAGCCGTTCTCTCCCTGTTCTTCCCGCACCCCTAAGCGATCCCCCGGGAATTC I E G R G I P A A V L L P V L L A A P ***				

FIG. 1

2/9




FIG. 2

3/9




FIG. 3a




FIG. 3b






FIG. 3b


4/9

*FIG. 4*


5/9

FIG. 5


PANEL 0




PANEL 1




PANEL 2




PANEL 3




PANEL 4



PANEL 5






PANEL 6



PANEL 7

6/9

*FIG. 6a**FIG. 6b**FIG. 6c*

7/9




FIG. 7a







FIG. 7b

8/9

*FIG. 8a**FIG. 8b*

9/9

*FIG. 9*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07189

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 38/00, 38/04; C07K 1/00; C12N 15/00, 15/64; C12P 21/06

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/324, 325, 326, 327, 328, 350; 435/320.1, 69.1, 91.4; 514/12, 13, 14, 15, 16

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

REGISTRY, CAPLUS, BIOSIS, MEDLINE, LIFESCI, EMBASE, USPATFULL, a-geneseq32, emb157, genbank110, n-geneseq32, pir58, swiss-prot35, sptrembl6, emb1-est-56, genbank-est109

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	ROJAS et al. Genetic engineering of proteins with cell membrane permeability. Nature Biotechnology. April 1998, Vol. 16, pages 370-372, relevant passages include the results, discussion and experimental protocol.	1-41, 44-54 and 55-59

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

25 MAY 1999

Date of mailing of the international search report

25 JUN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DEVESH SRIVASTAVA

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07189

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

530/324, 325, 326, 327, 328, 350; 435/320.1, 69.1, 91.4; 514/12, 13, 14, 15, 16

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.